arXiv:1304.6645v1 [cond-mat.stat-mech] 24 Apr 2013

The microscopic phase density functional approach to the construction of the kinetic and hydrodynamic description for the system of self-propelled particles

Oleksandr Chepizhko[1, 2] and Vladimir Kulinskii[1]

[1] *Department for Theoretical Physics, Odessa National University,*
Dvoryanskaya 2, 65026 Odessa, Ukraine

[2] *Laboratoire J.A. Dieudonné, Université de Nice Sophia Antipolis,*
UMR 7351 CNRS , Parc Valrose, F-06108 Nice Cedex 02, France

Abstract

We use the method of microscopic phase density to get general form of kinetic equation for the system of self-propelled particles with Vicsek-like alignment. The hydrodynamic equations are derived for the ordered phase and the ideal approximation is considered taking into account the mean-field force only. It is shown that the hydrodynamics of such ideal self-propelled fluid in main features demonstrates the dynamical transition from disordered initial state to the completely ordered motion. The problem of viscosity in hydrodynamic limit for the Vicsek-like model is discussed.

INTRODUCTION

The study of collective phenomena in systems of active agents is an important and fast developing field of statistical physics due to potential application to living matter dynamics [1, 2]. This phenomenon is widely observed in nature at different levels of organization including the synchronicity in insects behavior, bird flocking, traffic and complex social behavior [1, 3–5]. These systems are essentially nonequilibrium and their dynamics is not strictly Hamiltonian due to information exchange which includes not only positions but also the velocities.

One of the simplest models of the emergence of the coherent motion in active media is the (standard) Vicsek model (SVM) [6]. Basing on the simple rule it demonstrates the transition from disordered to the ordered state with varying the noise intensity. It has generated a series of publications [7–12] because of its simplicity and rich behavior. There are many questions that still need to be clarified for the model. One of the main open problem is the type of phase transition. Still it is not known exactly either it is continuous [13, 14] or discontinuous [9, 15]. It is still the subject of ongoing debate even on the "experimental" level of numerical simulation. The mean-field approximations of network model [16] and of the Vicsek model itself [17] demonstrate the dependence of the type of the transition on the type of noise. Approaching this problem theoretically leads to the formulation of proper kinetics equation which adequately reflects the basic physical mechanism of self-organization and justifies the hydrodynamic equations proposed previously from phenomenological reasonings [7, 18, 19]

Despite many results obtained in these approaches they lack the derivation of hydrodynamic equations which takes into account the specific features of SVM. The first attempt to do this was launched by Bertin et al. [11, 20] and Ihle [21, 22]. In first works kinetic equation was obtained within Boltzmann-like approach using two-particle collision integral. This gave the explicit expressions for the coefficients in hydrodynamic equations. But the results were quite different from numerical simulations, mainly because two-particle collisions are not sufficient to describe the model, and the usage of Boltzmann-like collision integral for active agents is not so evident, at least because the total momentum is not conserved. In the second works Liouville formalism was used. There the starting point was the equation for N-particle distribution function with subsequent factorization to one-particle function. The obtained kinetic equation took into account multi-particle collisions but neglected correlations between particles, which are very important in the VM. The shear viscosity terms appeared in all approaches. But until now no experimental (numerical) evidence of the viscous shear was given in the VM.

In this paper we use well-known method of microscopic phase density [23]. In the framework of this approach the kinetic equation is constructed directly on the basis of equations of motion without reference to N-particle distribution function. Also we work with velocities $\mathbf{v}$ which form such additive quantities like flow density and pressure. We use the angular variables only to calculate final values, unlike previous studies where the angular variables were used directly and this led to highly nonlinear functions and increased the complexity of calculations. To the best of our knowledge firstly such approach for the self-propelling system with Vicsek-type of interaction was used by P. Degond in [24, 25]. Although we also use the basic equation for the microscopic phase density functional as the starting point we proceed in physical approach in order to get the collision terms and hydrodynamic equations in a standard way (see [23, 26])

The paper is organized as follows. In Sec. I we derive the equation for microscopic phase density functional which leads to the formal kinetic equation with the corresponding collision terms. Then we obtain the hydrodynamical equations. In Sec. II we consider the hydrodynamical limit of ideal self-propelled Vicsek-like fluid. The question about the existence of the shear viscosity for such fluid is the subject of Sec. III. There we study the viscosity of this fluid from the theoretical point of view using the equations derived in previous sections. We postulate the problem of numerical simulations of Coutte flow for Vicsek-like model and discuss preliminary results. The analysis of the results and problems for future studies are given in concluding section.

I. THE EQUATION FOR THE MICROSCOPIC PHASE DENSITY FUNCTIONAL

The standard tool for the derivation of the hydrodynamic as well as kinetic equations for dynamical systems is the method of microscopic phase density functional [23]:

$$\mathcal{N}(x,t) = \sum_i \delta(x - x_i(t)) \tag{1}$$

where $x = (\mathbf{r}, \mathbf{v})$. It obeys the conservation law:

$$\partial_t \mathcal{N} + \mathbf{v}\, \partial_{\mathbf{r}} \mathcal{N} + \partial_{\mathbf{v}} \left(\dot{\mathbf{v}} \mathcal{N} \right) = 0\,. \tag{2}$$

The corresponding number density and the density flow are:

$$\rho^{(m)}(\mathbf{r},t) \; = \int \mathcal{N}(x,t)\, d\mathbf{v}\,, \tag{3}$$

$$\mathbf{j}^{(m)}(\mathbf{r},t) = \int \mathbf{v}\,\mathcal{N}(x,t)\, d\mathbf{v}\,, \tag{4}$$

where the superscript "m" stands for the "microscopic". We take the equation of motion for the Vicsek model [6] without noise term as following:

$$\frac{d\mathbf{v}_i}{dt} = \boldsymbol{\omega}_i \times \mathbf{v}_i \;, \tag{5}$$

which describes the fact that the kinetic energy of a particle is conserved. The original Vicsek algorithm [6] is based on the alignment of the particle velocity to the average direction of the neighbors:

$$\mathbf{w}_i = \frac{\sum_j \mathbf{v}_j K(\mathbf{r}_i - \mathbf{r}_j)}{|\sum_j \mathbf{v}_j K(\mathbf{r}_i - \mathbf{r}_j)|} \;, \tag{6}$$

where $K(\mathbf{r}_i - \mathbf{r}_j)$ is the "microsopic" summation kernel and therefore:

$$\boldsymbol{\omega}_i = \gamma\, \mathbf{v}_i \times \mathbf{w}_i \,. \tag{7}$$

Usually, the Heaviside step function is used for this purpose with the characteristic region of several average distance between the particles.

To avoid the difficulties in subsequent derivation connected with the non-additive structure of Eq. (6) and to make the transition to the low density limit $n \to 0$ trivial, we use another definition for $\mathbf{w}_i$:

$$\mathbf{w}_i = \sum_j \mathbf{v}_j K(\mathbf{r}_i - \mathbf{r}_j)\,, \tag{8}$$

i.e. the rate of alignment given by Eq. (8) is proportional to the local flux of the nearest neighbors. Otherwise using of Eq. (6) leads to the essential multiparticle collision terms (see e.g. [21]). These two options differ by the scalar factor γ. Therefore one can expect that despite the difference in microscopic equation of motion they lead in essential to the same macroscopic dynamics. Moreover, using Eq. (8) makes possible the comparison of the results of this approach with the Boltzmann like kinetic approach of Bertin et al. [20].

Because of its additive structure Eq. (8) can be expressed as the average of the MPDF:

$$\mathbf{j}_K^{(m)}(\mathbf{r},t) = \int dx'\, K(\mathbf{r}-\mathbf{r}')\mathbf{v}'\mathcal{N}(x',t) = \int K(\mathbf{r}-\mathbf{r}')\,\mathbf{j}^{(m)}(\mathbf{r}',t)\, d\mathbf{r}' \;. \tag{9}$$

Below we show that in such a case we get the equations similar to the usual case of molecular systems with pairwise interaction potential.

The equation (2) takes the form:

$$\partial_t \mathcal{N}(x,t) + \mathbf{v}\cdot\partial_{\mathbf{r}} \mathcal{N}(x,t) + \gamma \partial_{\mathbf{v}} \left(\int dx' \; (\, \mathbf{v} \times \mathbf{v}' \,) \times \mathbf{v}\, K(\mathbf{r}-\mathbf{r}') \mathcal{N}(x',t)\, \mathcal{N}(x,t) \right) = 0 \; . \quad (10)$$

From this equation it follows that the equation for one-particle distribution function $\overline{\mathcal{N}(x,t)} = n f_1(x,t)$ is:

$$\partial_t f_1(x,t) + \mathbf{v}\cdot\partial_{\mathbf{r}} f_1(x,t) + \frac{\gamma}{n} \partial_{\mathbf{v}} \left(\int K(\mathbf{r}-\mathbf{r}') \; (\, \mathbf{v} \times \mathbf{v}' \,) \times \mathbf{v}\, \overline{\mathcal{N}(x,t)\mathcal{N}(x',t)}\, dx' \right) = 0 \, , \quad (11)$$

where $\overline{\mathcal{N}(x,t)\mathcal{N}(x',t)}$ is connected with the pair distribution function f_2:

$$\overline{\mathcal{N}(x,t)\mathcal{N}(x',t)} = n\, \delta(x-x')\, f_1(x,t) + n^2\, f_2(x,x',t) \; . \quad (12)$$

Using common definition of the correlation function $g_2 = f_2 - f_1\, f_1$ we get:

$$\overline{(\delta\mathcal{N}\delta\mathcal{N})}_{x,x',t} = \overline{\mathcal{N}(x,t)\mathcal{N}(x',t)} - \overline{\mathcal{N}(x,t)}\; \overline{\mathcal{N}(x',t)} = n^2\, g_2(x,x',t) + n\, \delta(x-x')\, f_1(x,t) \; . \quad (13)$$

Thus Eq. (11) transforms into:

$$\left(\partial_t + \mathbf{v}\partial_{\mathbf{r}} + \partial_{\mathbf{v}} \mathbf{F}\right) f_1(x,t) = I(x,t) \; , \quad (14)$$

where $I(x,t)$ is the collision integral:

$$I(x,t) = \gamma\, n\, \partial_{\mathbf{v}} \left(\int \mathbf{v} \times (\mathbf{v} \times \mathbf{v}') \; K(\mathbf{r}-\mathbf{r}')\, g_2(x,x',t)\, dx' \right) \; . \quad (15)$$

Also we introduce the mean-field force $\mathbf{F}$ due to neighbors:

$$\mathbf{F}(x,t) = n\gamma \int \mathbf{v} \times (\, \mathbf{v}' \times \mathbf{v} \,) \; K(\mathbf{r}-\mathbf{r}')\, f_1(x',t)\, dx' = (\, \mathbf{v} \times \mathbf{j}_K \,) \times \mathbf{v} \, , \quad (16)$$

$$\mathbf{j}_K = \int K(\mathbf{r}-\mathbf{r}')\, \mathbf{v}'\, f_1(x',t)\, dx' = \int K(\mathbf{r}-\mathbf{r}')\, \mathbf{j}(\mathbf{r}',t)\, d\mathbf{r}' \; . \quad (17)$$

Using Eq. (14) the equations of motion for the basic physical quantities can be derived. The obvious conservation law for the number of particles is obtained from Eq. (14) by integration over $\mathbf{v}$:

$$\frac{\partial\rho}{\partial t} + \operatorname{div} \mathbf{j} = 0 \; , \quad (18)$$

where

$$\rho(\mathbf{r},t) = n \int f_1(x,t)\, d\mathbf{v} \, .$$

The equation for the hydrodynamic velocity field $\mathbf{u}$ has the form:

$$\frac{\partial\, (\, \rho\, u_i \,)}{\partial t} + \frac{\partial}{\partial x_j} (\, \rho\, u_i\, u_j \,) = -\frac{\partial P_{ij}}{\partial x_j} + \mathcal{F}_i + n \int \delta\, v_i\, I(x,t)\, d\mathbf{v} \; , \quad (19)$$

where the pressure tensor $\mathbf{P}$ can be decomposed into scalar and traceless components in a standard way:

$$P_{ij} = n \int \delta v_i \, \delta v_j \, f_1 \, d\mathbf{v} = p_0 \, \delta_{ij} + \pi_{ij} \ . \tag{20}$$

Here π_{ij} is the traceless part which can be identified with the viscosity stress tensor

$$\pi_{ij} = n \int \left(\delta v_i \, \delta v_j - \frac{\delta \mathbf{v}^2}{d} \delta_{ij} \right) f_1 \, d\mathbf{v} \ , \tag{21}$$

and

$$p_0 = n \int \frac{(\, \delta \mathbf{v} \,)^2}{2} f_1 \, d\mathbf{v} = \frac{\rho}{2} \left(1 - \mathbf{u}^2 \right) \ . \tag{22}$$

Also

$$\boldsymbol{\mathcal{F}}(\mathbf{r}, t) = n \int \mathbf{F} \, f_1(x, t) \, d\mathbf{v} \ , \tag{23}$$

is the mean force density. In the hydrodynamic limit the spatial scale of the variations for f_1 is much bigger than that of the kernel $K(\mathbf{r} - \mathbf{r}')$. Therefore the microscopic kernel should be put to δ-function and $\mathbf{j}_K \equiv \mathbf{j}$. Thus the force can be decomposed as following:

$$\boldsymbol{\mathcal{F}}(\mathbf{r}, t) = p_0 \, \mathbf{j} - \mathbf{j} . \boldsymbol{\pi} \ . \tag{24}$$

Note that in [24] the mean-field force was also obtained. It is proportional to the density gradient. But the representation in [24] did not provide clear distinction between this force and the dynamic term of the stress tensor because the authors used the field of unit vector $\boldsymbol{\Omega} = \mathbf{j}/|\mathbf{j}|$ of the local flux.

II. HYDRODYNAMIC LIMIT OF IDEAL SELF-PROPELLED VICSEK FLUID

Neglecting all fluctuational terms in Eq. (19) we get the equation of motion for an ideal self-propelling fluid as the continuous approximation of the Vicsek-like model:

$$\frac{d \, \mathbf{u}}{d \, t} = - \frac{\nabla \, p_0}{\rho} + \frac{\rho}{2} \left(1 - \mathbf{u}^2 \right) \mathbf{u} \tag{25}$$

where p_0 is given by Eq. (22). Although the first term in the right-hand side reminds the corresponding term of Euler equation for Newtonian ideal fluid. The second term as it follows from Eq. (24) is the mean force due to neighbors. Note that this term has the structure of Landau theory like terms introduced phenomenologically in [7]. One can expect that taking into account the collision terms will renormalize the unit coefficients of the corresponding



Figure 1. Density and velocity profiles with $v_0 = -1$.

power terms. The result Eq. (25) represents asymptotic form of the pressure terms in low noise limit. In the kinetic approach [20] the corresponding term has the form $(\mu - \xi\,\mathbf{u}^2)\,\mathbf{u}$ where μ, ξ are functions of density and the noise intensity. But the Boltzmann like approach of [20] is not applicable at low noise when $u \to 1$ since it is assumed that the density and the hydrodynamic velocity is low enough.

Using the obtained equations for the ideal spp-fluid it is natural to search for the running wave solutions of the form $\rho(\mathbf{r} - \mathbf{v}\,t)\,, \mathbf{u}(\mathbf{r} - \mathbf{v}\,t)$. Though Eq. (5) has sense only in dimensions more than 1 it is instructive to consider 1-dimensional case assuming the translational invariance along another directions. Here we give simple example of 1D solution $\rho(z)\,, u(z)$ with $z = x - v_0\,t$. The equations (18) and (25) can be integrated explicitly:

$$\rho(z) = \frac{C}{u(z) - v_0}\ , \tag{26}$$

where C is the constant of integration. The function $u(z)$ is determined by the equation:

$$\left(1 - 2v_0^2\right) \ln u(z) + v_0\,(v_0 - 1) \ln(1 - u(z)) + v_0\,(v_0 + 1) \ln(1 + u(z)) = C\,z\ . \tag{27}$$

Nonsingular solutions with $\rho(z)$ finite (see Eq. (26)) and $0 < u(z) < 1$ correspond to either $v_0 < 0$ or $v_0 > 1$. The asymptotic behavior at $t \to \infty$ of Eq. (27) depends on sign of v_0. If $v_0 < 0$ the order parameter field $u \to 0$ (see Fig. 1). If $v_0 > 1$ the transition to the completely ordered state $u \to 1$ takes place (see Fig. 2). Note that the solutions with



Figure 2. Density and velocity profiles with $v_0 = \sqrt{2}$.

$u \to 0$ at $t \to \infty$ correspond to the situation where the disordered region has higher density than the disordered one. This can be interpreted as the disappearance of unstable state. This is expected result because of the consideration of the ideal regime of low noise where all collision terms are neglected. Note that these solutions are similar to the switching wave solutions in an active bistable medium [27].

In Fig. 3 we also show the solution of equations (18), (25) under different initial states with periodic boundary conditions. It should be noted that these solutions asymptotically tend to the ordered state $u(x, t) \to 1$.

Obtained results indicate that the ideal fluid approximation still retains the basic features of initial self-propelling system despite neglecting the collision terms. It shows how different terms introduced from the heuristic arguments earlier (see e.g. [7, 19, 28]) arise.

III. ON THE VISCOSITY OF SELF-PROPELLED FLUID

In kinetic approach based on the Boltzman like arguments about the collision terms it is possible to get the equation of motion for the hydrodynamic fields [11, 20]. Yet the appearance of the viscous terms seems quite unexpected at least if the Vicsek-like model is assumed as the basis for the derivation of the kinetic equations. At the microscopic level the standard Vicsek model [29] is dissipativeless. Therefore one can expect the absence of



(a) $u(x,0) = 1 - \cos^8\left(\frac{\pi x}{2L}\right)$, $\rho(x,0) = 1 + \cos^4\left(\frac{\pi x}{2L}\right)$



(b) $u(x,0) = 1/4\,,\ \rho(x,0) = 1 + \cos^4\left(\frac{\pi x}{2L}\right)$

Figure 3. Dynamics of the velocity $u(x,t)$ and density $\rho(x,t)$ under different initial conditions.

viscosity at least in usual sense as the effect which decreases the momentum of the system.

Accurate derivation of the collision terms along with the account for the noise terms will be given in separate publication. Moreover different types of noise such as intrinsic [29] and the extrinsic one [9] apparently have different influence on the order-disorder transition at least within mean-field approximation [14, 16, 30].

Here we show that in the local equilibrium approximation the off-diagonal part of viscous tensor Eq. (21) vanishes. Indeed, as it follows from [17, 24, 31] in the local equilibrium approximation for the ordered phase the distribution function can be approximated as following:

$$f^{(0)}(\mathbf{r},\theta) = Ce^{a\cos\theta} \tag{28}$$

where C and a are some functions dependent on the local value $\mathbf{u}(\mathbf{r})$ of the order parameter. Obviously, it can serve as the interpolation between homogeneous distribution function $f = \frac{1}{2\pi}$ in the disordered state and the formal series representation of the distribution function near transition point to the ordered regime [20]:

$$f(\mathbf{r},\theta,t)=\frac{\rho(\mathbf{r},t)}{n}\left(\frac{1}{2\pi}+\frac{1}{\pi}u(\mathbf{r},t)\,\cos\theta+\dots\right)\ .$$

In such approximation the viscous tensor is:

$$\pi_{ij}^{(0)}=\int\left[\delta v_i\delta v_j-\delta_{ij}\frac{(\delta\mathbf{v})^2}{2}\right]f^{(0)}d\mathbf{v} \tag{29}$$

The off-diagonal components $i\neq j$ vanish

$$\pi_{12}=\pi_{21}=\int\delta v_1\delta v_2\,C\,e^{a\cos\theta}d\theta=C\int(\cos\theta-u)\sin\theta e^{a\cos\theta}d\theta=0 \tag{30}$$

according to the symmetry. The diagonal part of the tensor is determined by the quantity:

$$\Pi^{(0)}=\frac{1}{2}\int[\delta v_1^2-\delta v_2^2]f^{(0)}d\mathbf{v}=\frac{1}{2}\int[(\cos\theta-u)^2-\sin^2\theta]f^{(0)}d\theta\,. \tag{31}$$

which is not zero in general. Thus in this approximation the viscosity tensor is represented as:

$$\pi_{ij}^{(0)}=\Pi^{(0)}\begin{pmatrix}1&0\\0&-1\end{pmatrix} \tag{32}$$

Obtained results show that in this respect spp-fluid behaves differently from molecular liquids. For the ordinary fluid the local equilibrium is given by the Maxwell distribution and $\pi_{ij}^{(0)}=0$ due to basic properties of the collision integral [23]. We have shown that the shear viscosity is absent in the Vicsek-like fluid in local equilibrium approximation given by Eq. (28). Previous studies showed the viscosity terms in the hydrodynamic equations. So, this question needs to be resolved. The simplest way to check the existence of viscosity is a numerical simulation. To do this one can perform simulations of self-propelled particles system subjected to shear (as example [32]) and study the profile of the velocity field **u** that appears in such system. In the usual viscose liquid (Couette flow) linear profile (gradient) appears. Expected zero viscosity results in the absence of the nontrivial profile of the order parameter, except, probably, very thin layer close to the aligning boundary, caused by the specific mechanism of interaction.

We have performed a first simple check for a small system [33] and preliminary results show the absence of the order parameter profile. But the proper numerical answer to this problem is a question of our future study.

IV. DISCUSSION

In this paper we use the standard method of microscopic phase density to derive the general form of the kinetic equation for the Vicsek-like model. We obtain the form of the collision terms which in principle gives the possibility to obtain the closed kinetic equation for one-particle distribution function f_1. Such an approach is more closely connected with the equation of motion and could be easily modified to include different kind of velocity constraints and noise perturbation to dynamics. We have considered the simplest mean-field approximation and checked that the corresponding hydrodynamic equations have solutions displaying the transition to the ordered state. Note that we considered the rule of alignment to the local flux of the neighbors, but not to the average velocity as in original Vicsek model. By this the calculations are simplified because of the additivity and collision integrals depend on pair correlation function only. Although this simplification seems to be appropriate from the physical point of view, the question of mathematical equivalence between these two ways remains open and proper numerical study is required. Usage of the original Vicsek rule of alignment to the average velocity in principle lead to essentially many-particle collision terms. Thus in such an approach it is possible to consider in unified terms the kinetic equation obtained in Boltzmann-like derivation scheme [11, 20] and master equation approach [21] which lead to essentially many-particle collision terms.

Also we have considered important question about the viscosity of the Vicsek-like model. The calculation of the viscous tensor in the local approximation shows vanishing of the diagonal components of the tensor. We have checked this by looking at the velocity profile in a small test system. However proper numerical proofs of this result are required. The form of distribution function was taken as Boltzmann-like exponent from previous works [17, 31] though more rigorously it should be evaluated self-consistently on the basis of the kinetic equation. These and other questions will be considered in separate work.

ACKNOWLEDGMENTS

Authors are grateful to Professor Fernando Peruani and Dr. Francisco J. Sevilla for the fruitful discussion of the results.

[1] C. Castellano, S. Fortunato, and V. Loreto, Reviews of Modern Physics **81**, 591 (2009).

[2] T. Vicsek and A. Zafeiris, Physics Reports **517**, 71 (2012).

[3] D. Helbing, Rev. Mod. Phys. **73**, 1067 (2001).

[4] J. A. Acebrón, L. L. Bonilla, C. J. P. Vicente, F. Ritort, and R. Spigler, Reviews of Modern Physics **77**, 137 (2005).

[5] D. H. Kelley and N. T. Ouellette, Scientific Reports **3**, 1 (2013).

[6] T. Vicsek, A. Czirók, E. Ben-Jacob, I. Cohen, and O. Shochet, Physical Review Letters **75**, 1226 (1995).

[7] J. Toner and Y. Tu, Physical Review Letters **75**, 4326 (1995).

[8] M. Aldana and C. Huepe, Journal of Statistical Physics **112**, 135 (2003).

[9] G. Grégoire and H. Chaté, Phys. Rev. Lett. **92**, 025702 (2004).

[10] F. Peruani, A. Deutsch, and M. Bär, Physical Review E (Statistical, Nonlinear, and Soft Matter Physics) **74**,

[11] E. Bertin, M. Droz, and G. Grégoire, Physical review. E, Statistical, nonlinear, and soft matter physics **74** (2

[12] F. Cucker and S. Smale, Automatic Control, IEEE Transactions on **52**, 852 (2007).

[13] M. Nagy, I. Daruka, and T. Vicsek, Physica A: Statistical Mechanics and its Applications **373**, 445 (2007).

[14] M. Aldana, H. Larralde, and B. Vázquez, International Journal of Modern Physics B **23**, 3661 (2009), http://www.worldscientific.com/doi/pdf/10.1142/S0217979209053552.

[15] H. Chaté, F. Ginelli, G. Grégoire, and F. Raynaud, Phys. Rev. E **77**, 046113 (2008).

[16] M. Aldana, V. Dossetti, C. Huepe, V. M. Kenkre, and H. Larralde, Physical Review Letters **98**, 095702 (2007).

[17] A. A. Chepizhko and V. L. Kulinskii (AIP, 2009) pp. 25–33.

[18] J. Toner and Y. Tu, Phys. Rev. E **58**, 4828 (1998).

[19] V. L. Kulinskii, V. I. Ratushnaya, Zvelindovsky, and D. Bedeaux, Europhys. Lett. **71**, 207 (2005).

[20] E. Bertin, M. Droz, and G. Gregoire, Journal of Physics A: Mathematical and Theoretical **42**, 445001+ (200
arXiv:0907.4688.

[21] T. Ihle, Phys. Rev. E **83**, 030901 (2011).

[22] Y.-L. Chou, R. Wolfe, and T. Ihle, Phys. Rev. E **86**, 021120 (2012).

[23] Y. Klimontovich, *Statistical Physics* (Harwood Academic Publishers, NY, 1986).

[24] P. Degond and S. Motsch, Comptes Rendus Mathematique **345**, 555 (2007).

[25] P. Degond and S. Motsch, Journal of Statistical Physics , 2008.

[26] E. M. Lifshitz and L. P. Pitaevskii, *Course of Theoretical Physics, Volume X: Physical Kinetics* (Butterworth-Heinemann, 1981).

[27] A. S. Mikhailov and A. Y. Loskutov, *Foundations Of Synergetics II - Complex Patterns* (Springer-Verlag, 1996).

[28] Y. Tu, J. Toner, and M. Ulm, Physical Review Letters **80**, 4819 (1998).

[29] A. Czirók, E. Ben-Jacob, I. Cohen, and T. Vicsek, Phys. Rev. E **54**, 1791 (1996).

[30] A. Chepizhko and V. Kulinskii, Physica A: Statistical Mechanics and its Applications **389**, 5347 (2010).

[31] Z. Csahók and T. Vicsek, Physical Review E **52**, 5297+ (1995).

[32] D. M. E. O. Gustavo P. Saracco, Giuseppe Gonnella, Central European Journal of Physics **10**, 1109 (2012).

[33] V. Kulisnkii and A. Chepizhko, "Self-propelling particles as a new challenge for nonequilibrium statistical me
presneted at 6th International Workshop on Nonequilibrium Thermodynamics and 3rd Lars Onsager Symposium IWNET 2012, August 19-24, Roros, Norway (2012).